



HAL
open science

FINITE CONVERGENT PRESENTATIONS OF PLACTIC MONOIDS FOR SEMISIMPLE LIE ALGEBRAS

Nohra Hage

► **To cite this version:**

Nohra Hage. FINITE CONVERGENT PRESENTATIONS OF PLACTIC MONOIDS FOR SEMISIMPLE LIE ALGEBRAS. 2015. ujm-01248274

HAL Id: ujm-01248274

<https://ujm.hal.science/ujm-01248274>

Preprint submitted on 24 Dec 2015

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

FINITE CONVERGENT PRESENTATIONS OF PLACTIC MONOIDS FOR SEMISIMPLE LIE ALGEBRAS ¹

NOHRA HAGE

Abstract – We study rewriting properties of the column presentation of plactic monoid for any semisimple Lie algebra such as termination and confluence. Littelmann described this presentation using L-S paths generators. Thanks to the shapes of tableaux, we show that this presentation is finite and convergent. We obtain as a corollary that plactic monoids for any semisimple Lie algebra satisfy homological finiteness properties.

Keywords – Plactic algebra; Littelmann path model; standard tableau; convergent presentations.
M.S.C. 2010 – 16S15, 68Q42, 20M05, 68R05, 06B15

1. INTRODUCTION

Using his path model, Littelmann defined in [Lit96] a plactic algebra for any semisimple Lie algebra. As a consequence, he gave some presentations by generators and relations of the plactic algebra of types A, B, C, D and G_2 . Using a case-by-case analysis, the plactic congruence can be also defined using Kashiwara's theory of crystal bases [DJM90, Kas91, KN94, Kas95]. Lascoux, Leclerc and Thibon presented in [LLT95] the plactic monoid of type A using the theory of crystal bases and gave a presentation of the plactic monoid of type C without proof. Shortly after that, Lecouvey in [Lec02] and Baker in [Bak00] described independently the plactic monoid of type C using also Kashiwara's theory of crystal bases. Plactic monoids of types B, D and G_2 were introduced by Lecouvey, see [Lec07].

The plactic monoid of rank n introduced by Lascoux and Schützenberger in [LS81], corresponds to the representations of the general linear Lie algebra of n by n matrices. This Lie algebra is of type A, so the corresponding plactic monoid is known as the plactic monoid of type A, and denoted by $\mathbf{P}_n(A)$.

The monoid $\mathbf{P}_n(A)$ has found several applications in algebraic combinatorics and representation theory [LS81, LLT95, Ful97, Lot02]. Similarly, plactic monoids for any types have many applications like the combinatorial description of the Kostka-Foulkes polynomials which arise as entries of the character table of the finite reductive groups [Lec06].

More recently, plactic monoids were investigated by rewriting methods. In [KO14], Kubat and Okninski showed that for $n > 3$, there is no finite completion of the presentation of the monoid $\mathbf{P}_n(A)$ with the Knuth generators. Bokut et al. in [BCCL15] and Cain et al. in [CGM15] constructed independently finite convergent presentation of the monoid $\mathbf{P}_n(A)$ by adding columns generators. This presentation is called the *column presentation*. Having a finite convergent presentation of the monoid $\mathbf{P}_n(A)$ had many consequences. In particular, the column presentation was used by Lopatkin in [Lop14] to construct Anick's resolution for the monoid $\mathbf{P}_n(A)$ and by the author and Malbos in [HM15] to construct coherent

¹THIS WORK IS PARTIALLY SUPPORTED BY THE FRENCH NATIONAL RESEARCH AGENCY, ANR-13-BS02-0005-02.

presentations of it. Note that coherent presentation extends the notion of a presentation of the monoid by homotopy generators taking into account the relations among the relations. Using Kashiwara's theory of crystal bases, the author generalized the column presentation for type A and constructed a finite convergent presentation of plactic monoid of type C by adding admissible column generators [Hag14]. A bit later, a similar column presentation was obtained by Cain et al. for plactic monoids of types B, C, D and G_2 [CGM14].

In the present work, we consider plactic monoids for any semisimple Lie algebra. The column presentation of these monoids was introduced by Littelmann in [Lit96] using L-S paths. An L-S path corresponds to a column for type A, and to an admissible column for types C, B, D and G_2 in the Lecouvey sense, see [Lec07]. We study the column presentation using rewriting methods. For this, we consider a rewriting system where the generating set contains the finite set of L-S paths. The right-hand sides of rewriting rules are standard tableaux. The rewriting system rewrites two L-S paths that their concatenation do not form a standard tableau to their corresponding standard tableau form. Using the shapes of tableaux, we show that this presentation is finite and convergent. As a consequence, we deduce that plactic monoids for any types satisfy some homological finiteness properties.

Note that the convergent column presentation of plactic monoids for any semisimple Lie algebra given in this paper using L-S paths coincides with the presentations constructed for type A in [CGM15, BCCL15], for type C in [Hag14] and for types B, C, D and G_2 in [CGM14].

The paper is organised as follows. In Section 2, we recall the definitions of paths, root operators and L-S paths. After we present the definition and some properties of standard tableaux as defined by Littelmann and we recall the definition of the plactic algebra for any semisimple Lie algebra. In Section 3, we recall some rewriting properties of the presentations of monoids in term of polygraphs. After, we show that the column presentation of the plactic monoid for any semisimple Lie algebra is finite and convergent.

2. PLACTIC ALGEBRA

2.1. L-S Paths

In this section, we recall the definitions and properties of paths, root operators and L-S paths. We refer the reader to [Lit94, Lit95, Lit96] for a full introduction.

2.1.1. Root system. Let \mathfrak{g} be a semisimple Lie algebra. Let V be the finite vector space with standard inner product $\langle \cdot, \cdot \rangle$ spanned by $\Phi \subset V \setminus \{0\}$ the root system of \mathfrak{g} . Let Φ^+ be the set of its positive roots. A positive root α in Φ^+ is *simple*, if it can not be written as $\alpha_1 + \alpha_2$, where $\alpha_1, \alpha_2 \in \Phi^+$. For a root α , define by $\alpha^\vee := \frac{2\alpha}{\langle \alpha, \alpha \rangle}$ its *coroot*. Denote by

$$X = \{ v \in V \mid \langle v, \alpha_i^\vee \rangle \in \mathbb{Z}, \text{ for all } i \}$$


the weight lattice of the Lie algebra \mathfrak{g} and set $X_{\mathbb{R}} := X \otimes_{\mathbb{Z}} \mathbb{R}$. A *fundamental weight* ω_i in X satisfy $\langle \omega_i, \alpha_j^\vee \rangle = \delta_{ij}$, for all i and j . The set of dominant weights is

$$X^+ = \{ \lambda \in X \mid \langle \lambda, \alpha_i^\vee \rangle \geq 0, \text{ for all } i \}$$


and the dominant chamber is

$$X_{\mathbb{R}}^+ = \{ \lambda \in X_{\mathbb{R}} \mid \langle \lambda, \alpha_i^\vee \rangle \geq 0, \text{ for all } i \}.$$

2.1.2. Example. Let $\mathfrak{g} = \mathfrak{sl}_3$. Consider $V = \{ (x_1, x_2, x_3) \in \mathbb{R}^3 \mid x_1 + x_2 + x_3 = 0 \}$ and let $\{\varepsilon_1, \varepsilon_2, \varepsilon_3\}$ be the canonical basis of \mathbb{R}^3 . The simple roots of \mathfrak{g} are $\alpha_1 = \varepsilon_1 - \varepsilon_2$ and $\alpha_2 = \varepsilon_2 - \varepsilon_3$. Its fundamental weights are $\omega_1 = \varepsilon_1$ and $\omega_2 = \varepsilon_1 + \varepsilon_2$ (we still denote by ε_i the projection of ε_i into V). An example of dominant weight is $\omega_1 + \omega_2$. The dominant chamber is the hatched area on the following figure:



2.1.3. Example. Let $\mathfrak{g} = \mathfrak{sp}_4$. Consider $V = \mathbb{R}^2$ with its canonical basis $\{\varepsilon_1, \varepsilon_2\}$. The simple roots are $\alpha_1 = \varepsilon_1 - \varepsilon_2$ and $\alpha_2 = 2\varepsilon_2$. The fundamental weights are $\omega_1 = \varepsilon_1$ and $\omega_2 = \varepsilon_1 + \varepsilon_2$. An example of dominant weight is $\omega_1 + \omega_2$. The dominant chamber is the hatched area on the following figure:



For more informations, we refer the reader to [Bou68, Hum78].

2.1.4. Paths. A *path* is a piecewise linear, continuous map $\pi : [0, 1] \rightarrow X_{\mathbb{R}}$. We will consider paths up to a reparametrization, that is, the path π is equal to any path $\pi \circ \varphi$, where $\varphi : [0, 1] \rightarrow [0, 1]$ is a piecewise linear, nondecreasing, surjective, continuous map. The *weight* of a path π is $\text{wt}(\pi) = \pi(1)$. For example, for the Lie algebra $\mathfrak{g} = \mathfrak{sl}_n$, the paths $\pi_{\varepsilon_i} : t \mapsto t\varepsilon_i$ are of weight ε_i , for $i = 1, \dots, n$. Denote by

$$\Pi = \{ \pi : [0, 1] \rightarrow X_{\mathbb{R}} \mid \pi(0) = 0 \text{ and } \pi(1) \in X \}$$

the set of all paths with sources 0 such that their weight lies in X . Given two paths π_1 and π_2 in Π , the *concatenation* of π_1 and π_2 , denoted by $\pi := \pi_1 \star \pi_2$, is defined by:


$$\pi(t) := \begin{cases} \pi_1(2t) & \text{for } 0 \leq t \leq \frac{1}{2} \\ \pi_1(1) + \pi_2(2t - 1) & \text{for } \frac{1}{2} \leq t \leq 1 \end{cases}$$

2. Plactic algebra

Denote by $\mathbb{Z}\Pi$ the *algebra of paths* defined as the free \mathbb{Z} -module with basis Π whose product is given by the concatenation of paths and where the unity is the trivial path

$$\begin{aligned} \theta : [0, 1] &\longrightarrow X_{\mathbb{R}} \\ t &\longmapsto 0. \end{aligned}$$

2.1.5. Example. Let $\mathfrak{g} = \mathfrak{sl}_3$. Consider the paths $\pi_1 : t \mapsto t\varepsilon_1$ and $\pi_2 : t \mapsto t\varepsilon_2$. The path $\pi_1 \star \pi_2$ is the path on the following figure:



2.1.6. Root operators. For each simple root α , one defines root operators $e_\alpha, f_\alpha : \Pi \rightarrow \Pi \cup \{0\}$ as follows. Every path π in Π is cutted into three parts, *i.e.*, $\pi = \pi_1 \star \pi_2 \star \pi_3$. Then the new path $e_\alpha(\pi)$ or $f_\alpha(\pi)$ is either equal to 0 or $\pi_1 \star s_\alpha(\pi_2) \star \pi_3$, where s_α denote the simple reflection with respect to the root α . In other words, consider the function

$$\begin{aligned} h_\alpha : [0, 1] &\longrightarrow \mathbb{R} \\ t &\longmapsto \langle \pi(t), \alpha^\vee \rangle \end{aligned}$$

Let $Q := \min(\text{Im}(h_\alpha) \cap \mathbb{Z})$ be the minimum attained by h_α . If $Q = 0$, define $e_\alpha(\pi) = 0$. If $Q < 0$, let

$$t_1 = \min\{ t \in [0, 1] \mid h_\alpha(t) = Q \}$$

and

$$t_0 = \max\{ t < t_1 \mid h_\alpha(t) = Q + 1 \}.$$

Denote by π_1, π_2 and π_3 the paths defined by

$$\begin{aligned} \pi_1(t) &:= \pi(tt_0) \\ \pi_2(t) &:= \pi(t_0 + t(t_1 - t_0)) - \pi(t_0) \\ \pi_3(t) &:= \pi(t_1 + t(1 - t_1)) - \pi(t_1), \text{ for } t \in [0, 1]. \end{aligned}$$

By definition of the π_i , we have $\pi = \pi_1 \star \pi_2 \star \pi_3$. Then $e_\alpha(\pi) = \pi_1 \star s_\alpha(\pi_2) \star \pi_3$.

Similarly, one can define the operator f_α . Let

$$p = \max\{ t \in [0, 1] \mid h_\alpha(t) = Q \}.$$

Denote by P the integral part of $h_\alpha(1) - Q$. If $P = 0$, define $f_\alpha(\pi) = 0$. If $P > 0$, let $x > p$ such that

$$h_\alpha(x) = Q + 1 \text{ and } Q < h_\alpha(x) < Q + 1, \text{ for } p < t < x.$$

Denote by π_1, π_2 and π_3 the paths defined by

$$\begin{aligned}\pi_1(t) &:= \pi(tp) \\ \pi_2(t) &:= \pi(p + t(x - p)) - \pi(p) \\ \pi_3(t) &:= \pi(x + t(1 - x)) - \pi(x), \text{ for } t \in [0, 1].\end{aligned}$$

By definition of the π_i , we have $\pi = \pi_1 \star \pi_2 \star \pi_3$. Then $f_\alpha(\pi) = \pi_1 \star s_\alpha(\pi_2) \star \pi_3$.

These operators preserve the length of the paths. We have also that if $f_\alpha(\pi) = \pi' \neq 0$ then $e_\alpha(\pi') = \pi \neq 0$ and $\text{wt}(f_\alpha(\pi)) = \text{wt}(e_\alpha(\pi)) - \alpha$.


For all simple root α , let \mathcal{A} be the subalgebra of $\text{End}_{\mathbb{Z}}(\mathbb{Z}\Pi)$ generated by the root operators f_α and e_α . Define Π^+ to be the set of paths π such that the image is contained in $X_{\mathbb{R}}^+$ and denote by M_π the \mathcal{A} -module $\mathcal{A}\pi$. Let B_π be the \mathbb{Z} -basis $M_\pi \cap \Pi$ of M_π . In other words, for a finite set I , we have

$$B_\pi = \{ f_{i_1} \dots f_{i_r}(\pi) \mid \pi \in \Pi^+ \text{ and } i_1, \dots, i_r \in I \}.$$

For a dominant weight λ , consider the path $\pi_\lambda(t) = t\lambda$ that connects the origin with λ by a straight line. Denote by M_λ the \mathcal{A} -module $\mathcal{A}\pi_\lambda$ generated by the path π_λ . In addition, the \mathbb{Z} -module M_λ has for a basis the set B_{π_λ} consisting of all paths in M_λ .

Given two paths π and π' in Π^+ , if $\pi(1) = \pi'(1)$, then the \mathcal{A} -modules M_π and $M_{\pi'}$ are isomorphic. For π in Π^+ , let η in M_π be an arbitrary path. The minimum of the function $h_\alpha(t) = \langle \pi(t), \alpha^\vee \rangle$ is an integer for all simple roots. In addition, we have $e_\alpha(\eta) = 0$ if and only if $\eta = \pi$.

2.1.7. Example. Let $\mathfrak{g} = \mathfrak{sl}_3$ and let α_1, α_2 be its simple roots and ω_1, ω_2 be its fundamental weights. For $\lambda = \omega_1 + \omega_2$, consider the path $\pi_\lambda : t \mapsto t\lambda$. Let us compute B_{π_λ} . We have



2. Plactic algebra


where

$$\begin{aligned}
 \pi_\lambda &: t \longrightarrow t\lambda \\
 f_{\alpha_1}(\pi_\lambda) &=: \pi_2 : t \longrightarrow t\alpha_2 \\
 f_{\alpha_2}(\pi_\lambda) &=: \pi_3 : t \longrightarrow t\alpha_1 \\
 f_{\alpha_2}(\pi_2) &=: \pi_4 : t \longrightarrow \begin{cases} -t\alpha_2 & \text{for } 0 \leq t \leq \frac{1}{2} \\ (t-1)\alpha_2 & \text{for } \frac{1}{2} \leq t \leq 1 \end{cases} \\
 f_{\alpha_1}(\pi_3) &=: \pi_5 : t \longrightarrow \begin{cases} -t\alpha_1 & \text{for } 0 \leq t \leq \frac{1}{2} \\ (t-1)\alpha_1 & \text{for } \frac{1}{2} \leq t \leq 1 \end{cases} \\
 f_{\alpha_2}(\pi_4) &=: \pi_6 : t \longrightarrow -t\alpha_2 \\
 f_{\alpha_1}(\pi_5) &=: \pi_7 : t \longrightarrow -t\alpha_1 \\
 f_{\alpha_2}(\pi_7) = f_{\alpha_1}(\pi_6) &=: \pi_8 : t \longrightarrow -t\lambda
 \end{aligned}$$

Hence we obtain $B_{\pi_\lambda} = \{\pi_\lambda, \pi_2, \pi_3, \pi_4, \pi_5, \pi_6, \pi_7, \pi_8\}$.

2.1.8. Crystal graphs. For π in Π^+ , let $\mathcal{G}(\pi)$ be the oriented graph with set of vertices B_π and an arrow $\pi \xrightarrow{i} \pi'$ means that $f_{\alpha_i}(\pi) = \pi'$ and $e_{\alpha_i}(\pi') = \pi$. The graph $\mathcal{G}(\pi)$ is also called a *crystal graph*, see [Kas95].

For example, consider the semisimple Lie algebra $\mathfrak{g} = \mathfrak{sl}_3$. For $\lambda = \omega_1 + \omega_2$, the elements of B_{π_λ} obtained in Example 2.1.7 can be represented in the following crystal graph $\mathcal{G}(B_{\pi_\lambda})$



2.1.9. L-S paths. For a dominant weight λ , the *Lakshmibai-Seshadri paths*, *L-S paths* for short, of shape λ are the paths π of the form

$$\pi = f_{\alpha_1} \circ \dots \circ f_{\alpha_s}(\pi_\lambda)$$

where $\alpha_1, \dots, \alpha_s$ are simple roots of a semisimple Lie algebra \mathfrak{g} . That is, these paths are all the elements of B_{π_λ} .

2.1.10. Example. Consider $\mathfrak{g} = \mathfrak{sl}_3$ and let ω_1 and ω_2 be its fundamental weights. Let $\lambda = \omega_1 + \omega_2$, the L-S paths of shape λ are all the elements of B_{π_λ} as obtained in Example 2.1.7. The L-S paths of shape ω_2 are the elements of $B_{\pi_{\omega_2}}$, with


$$B_{\pi_{\omega_2}} = \{ \pi_{\varepsilon_1 + \varepsilon_2}, \pi_{\varepsilon_1 + \varepsilon_3}, \pi_{\varepsilon_2 + \varepsilon_3} \}.$$

For the Lie algebra \mathfrak{sl}_n , the L-S paths of shape ω_1 are the paths π_{ε_i} , for $i = 1, \dots, n + 1$.

2.2. Tableaux

In this section, we recall the definition of standard tableaux as defined by Littelmann in [Lit96]. Fix an enumeration $\omega_1, \dots, \omega_n$ of the fundamental weights of a semisimple Lie algebra \mathfrak{g} .

2.2.1. Young diagram. A *Young diagram* is a finite collection of boxes, arranged in left-justified rows such that the length of each row is bigger or equal to the one below it. Note that the rows are enumerated from top to bottom and the columns from the right to the left. For example, the Young diagram with six boxes in the first row, four boxes in the second boxes and one box in the third row is the following



2.2.2. L-S monomials. Let $\lambda = \lambda_1 + \dots + \lambda_k$, where $\lambda_1, \dots, \lambda_k$ are dominant weights. If for all $i = 1, \dots, k$, π_i is an L-S path of shape λ_i , then the monomial $m = \pi_1 \star \dots \star \pi_k$ is called an *L-S monomial of shape $\underline{\lambda} = (\lambda_1, \dots, \lambda_k)$* .

2.2.3. Young tableau. A *Young tableau of shape $\lambda = a_1\omega_1 + \dots + a_n\omega_n$* is an L-S monomial

$$\star_{\substack{1 \leq i \leq n \\ 1 \leq j \leq n}} \pi_{1, \omega_j} \star \dots \star \pi_{a_i, \omega_i}$$


where π_{i, ω_j} is an L-S path of shape ω_j , for $1 \leq i \leq n$. That is, the first a_1 paths are of shape ω_1 , the next a_2 are of shape ω_2, \dots , the final a_n paths are of shape ω_n .

2.2.4. Standard tableau. Let m be a Young tableau of shape $\lambda = a_1\omega_1 + \dots + a_n\omega_n$. The Young tableau m is called *standard of shape λ* if $m \in \mathcal{A}(\pi_{\omega_1} \star \dots \star \pi_{\omega_n})$.

2.2.5. Example. For type A_n , consider the ordered alphabet $\mathcal{A}_{n+1} = \{1 < \dots < n+1\}$. For a dominant weight $\lambda = a_1\omega_1 + \dots + a_n\omega_n$, set $p_i = a_i + \dots + a_n$. Consider the Young diagram with p_1 boxes in the first row, p_2 boxes in the second row, etc. A Young tableau of shape λ and type A is a filling of the boxes of this Young diagram with elements of the alphabet \mathcal{A}_{n+1} such that the entries are strictly increasing in the column from top to bottom.

A standard tableau of shape $\lambda = \omega_1 + \dots + \omega_k$ is a Young tableau of shape λ and type A such that the entries are weakly increasing in the rows from left to right. In other words all the standard tableaux of shape λ are the elements of $B_{\pi_{\omega_1} \star \dots \star \pi_{\omega_k}}$, where the L-S monomial $\pi_{\omega_1} \star \dots \star \pi_{\omega_k}$ corresponds to the Young tableau with only 1's in the first row, 2's in the second row, \dots , k 's in the k -th row. Note that the standard tableaux of type A are also called *semistandard tableaux* in [Ful97].

For type A_3 , the following Young tableau is standard of shape $\omega_1 + 2\omega_2 + \omega_3$



2.2.6. Example. For type C_n , consider the ordered alphabet $\mathcal{C}_n = \{1 < \dots < n < \bar{n} < \dots < \bar{2} < \bar{1}\}$. For a dominant weight $\lambda = a_1\omega_1 + \dots + a_n\omega_n$, set $p_i = a_1 + 2a_2 + \dots + 2a_n$ and for $i \geq 2$

3. Finite convergent presentation of plactic monoids

set $p_i = 2a_i + \dots + 2a_n$. Consider the Young diagram with p_1 boxes in the first row, p_2 boxes in the second row, etc. A Young tableau of shape λ and type C is a filling of the boxes of this Young diagram with elements of the alphabet \mathcal{C}_n such that the entries are strictly increasing in the column from top to bottom, but \bar{i} and i are never entries in the same column. In addition, for each pair of columns $(C_{a_1+2j-1}, C_{a_1+2j})$, $j = 1, \dots, a_2 + \dots + a_n$, either these columns are equal or the column C_{a_1+2j} is obtained from C_{a_1+2j-1} by exchanging an even number of times an entry k , $1 \leq k \leq \bar{1}$, in C_{a_1+2j-1} by \bar{k} , see [GL12, Section 4.2].

A standard tableau of shape $\lambda = \omega_1 + \dots + \omega_k$ is a Young tableau of shape λ and type C such that the entries are weakly increasing in the rows from left to right. Note that the standard tableaux of type C are also called *symplectic tableaux* in [Lec02].

For type C_3 , the following Young tableau is a standard tableau of shape $\omega_1 + 2\omega_2$

1	1	1	2	$\bar{3}$
2	2	$\bar{3}$	$\bar{3}$	
3	3	2	$\bar{1}$	

2.3. Plactic algebra for any semisimple Lie algebra

Let us recall the definition of the plactic algebra for any semisimple Lie algebra and we refer the reader to [Lit96] for more details.

Let $\mathbb{Z}\Pi_0$ be the \mathcal{A} -submodule $\mathcal{A}\Pi^+$ of $\mathbb{Z}\Pi$ generated by the paths in Π^+ . For two paths π_1 and π_2 in $\mathbb{Z}\Pi_0$, denote by π_1^+ and π_2^+ the unique paths in Π^+ such that $\pi_1 \in M_{\pi_1^+}$ and $\pi_2 \in M_{\pi_2^+}$. One can define a relation \sim on $\mathbb{Z}\Pi_0$ by: $\pi_1 \sim \pi_2$ if, and only if, $\text{wt}(\pi_1^+) = \text{wt}(\pi_2^+)$ and $\psi(\pi_1) = \pi_2$ under the isomorphism $\psi : M_{\pi_1^+} \rightarrow M_{\pi_2^+}$. The *plactic algebra* for \mathfrak{g} is the quotient

$$\mathbb{Z}\mathcal{P} := \mathbb{Z}\Pi_0 / \sim.$$

For $\pi \in \mathbb{Z}\Pi_0$, we denote by $[\pi] \in \mathbb{Z}\mathcal{P}$ its equivalence class. The classes $[m]$ of standard Young tableaux form a basis of the plactic algebra $\mathbb{Z}\mathcal{P}$, see [Lit96, Theorem 7.1].

2.3.1. Example ([Lit96, Theorem C]). For type A, consider the alphabet $\{1, \dots, n\}$. The plactic congruence coincides with the congruence generated by the following families of relations on the word algebra $\mathbb{Z}\{1, \dots, n\}$:

- (a) $xzy = zxy$ for $1 \leq x < y \leq z \leq n$.
- (b) $yxz = yzx$ for $1 \leq x \leq y < z \leq n$.
- (c) $12 \dots n$ is the trivial word.

3. FINITE CONVERGENT PRESENTATION OF PLACTIC MONOIDS

3.1. Rewriting properties of 2-polygraphs

In this section, we recall some rewriting properties of the presentations of monoids. These presentations are studied in terms of polygraphs. For more informations, we refer the reader to [GM14].

A *2-polygraph* (with only one 0-cell) is a pair $\Sigma = (\Sigma_1, \Sigma_2)$ where Σ_1 is the set of generating 1-cells and Σ_2 is the set of generating 2-cells, that is rules of the form $u \Rightarrow v$, with u and v are words in the free monoid Σ_1^* . A monoid M is presented by a 2-polygraph Σ if M is isomorphic to the quotient of the free monoid Σ_1^* by the congruence generated by Σ_2 . A 2-polygraph Σ is *finite* if Σ_1 and Σ_2 are finite. A *rewriting step* of Σ is a 2-cell of the form $w\varphi w' : wuww' \Longrightarrow wvww'$, where $\varphi : u \Longrightarrow v$ is a 2-cell in Σ_2 and w and w' are words of Σ_1^* . A *rewriting sequence* of Σ is a finite or infinite sequence of rewriting steps. We say that u rewrites into v if Σ has a nonempty rewriting sequence from u to v . A word of Σ_1^* is a *normal form* if Σ has no rewriting step with source u . A normal form of u is a word v of Σ_1^* that is a normal form and such that u rewrites into v . We say that Σ *terminates* if it has no infinite rewriting sequences. We say that Σ is *confluent* if for any words u, u' and u'' of Σ_1^* , such that u rewrites into u' and u'' , there exists a word v in Σ_1^* such that u' and u'' rewrite into v . We say that Σ is *convergent* if it terminates and it is confluent. Note that a terminating 2-polygraph is convergent if every word admits a unique normal form.

3.2. Column presentation

Let \mathfrak{g} be a semisimple Lie algebra and let $\omega_1, \dots, \omega_n$ be its fundamental weights. Let \mathbb{B}_i be the set of L-S paths of shape ω_i and $\mathbb{B} = \cup_{i=1}^n \mathbb{B}_i$. Denote by \mathbb{B}^* the free monoid over \mathbb{B} . A word on \mathbb{B}^* is a concatenation of L-S paths.

For every L-S paths c_1 and c_2 in \mathbb{B} such that $c_1 \star c_2$ is not a standard tableau, we define a 2-cell

$$c_1 \star c_2 \xrightarrow{\gamma_{c_1, c_2}} T$$

where T is the unique standard tableau such that $[T] = [c_1 \star c_2]$.

Let denote by $\text{Path}(n)$ the 2-polygraph whose set of 1-cells is \mathbb{B} and the set of 2-cells is

$$\text{Path}_2(n) = \{ c_1 \star c_2 \xrightarrow{\gamma_{c_1, c_2}} T \mid c_1, c_2 \in \mathbb{B} \text{ and } c_1 \star c_2 \text{ is not a standard tableau} \}.$$

This presentation is called the *column presentation*. It is a presentation of the plactic monoid for any semisimple Lie algebra [Lit96, Theorem B]. Let us prove that it is terminating and confluent.

3.2.1. Order on the tableaux. Consider the partial order \leq of dominant weights defined by

$$\lambda_1 \leq \lambda_2 \quad \text{if and only if} \quad \lambda_1 - \lambda_2 \in \mathbb{N}\Phi^+$$

where λ_1 and λ_2 are dominant weights. That is, $\lambda_1 - \lambda_2$ is a nonnegative integral sum of positive roots. Using this order, one can find for each dominant weight, a finite number of dominant weights that are smaller than it, then the partial order \leq is a well-founded order.

Let us define an order \preceq on the tableaux of \mathbb{B}^* as follows. For two tableaux m and m' of shape λ and λ' respectively, we have

$$m \preceq m' \quad \text{if and only if} \quad \lambda \leq \lambda'.$$

3.2.2. Remark. In the proof of Theorem B in [Lit96], Littelmann showed that for every tableau m of shape λ that is not standard, the shape of the standard tableau m' such that $[m] = [m']$ is strictly smaller than λ .

3. Finite convergent presentation of plactic monoids

3.2.3. Lemma. *The 2-polygraph $\text{Path}(n)$ is terminating.*

Proof. Let us show that the 2-polygraph $\text{Path}(n)$ is compatible with the order \preceq . We have to prove that if $h \implies h'$, then $h' \prec h$, where h is not a standard tableau and h' is a standard tableau of shape λ' . There are two cases depending on whether or not h is a tableau.

Suppose that h is a tableau of shape λ that it is not standard, then by Remark 3.2.2 we have $\lambda' < \lambda$. Thus we obtain that $h' \prec h$.

Suppose that h is an L-S monomial of shape $\underline{\lambda}$ that it is not a tableau. By decomposing each L-S path of shape ω_i , for $i = 1, \dots, n$, in h into concatenation of L-S paths of shape ω_1 , we transform the L-S monomial h into a tableau of shape $k\omega_1$, with $k \in \mathbb{N}$. Again by Remark 3.2.2, we have $\lambda' < k\omega_1$. Thus we obtain that $h' \prec h$.

Hence, rewriting an L-S monomial that is not a standard tableau always decreases it with respect to the order \preceq . Since every application of a 2-cell of $\text{Path}(n)$ yields a \prec -preceding tableau, it follows that any sequence of rewriting using $\text{Path}(n)$ must terminate. \square

3.2.4. Lemma. *The 2-polygraph $\text{Path}(n)$ is confluent.*

Proof. Let $m \in \mathbb{B}^*$ and T, T' be two normal forms such that $m \implies T$ and $m \implies T'$. It is sufficient to prove that $T = T'$. Suppose $T = c_1 \star \dots \star c_k$, where the L-S monomial $c_1 \star \dots \star c_k$ is a standard Young tableau such that $[m] = [c_1 \star \dots \star c_k]$. Similarly, $T' = c'_1 \star \dots \star c'_l$ where the L-S monomial $c'_1 \star \dots \star c'_l$ is a standard Young tableau such that $[m] = [c'_1 \star \dots \star c'_l]$. Since $[m] = [c_1 \star \dots \star c_k] = [c'_1 \star \dots \star c'_l]$ and the standard tableaux form a cross-section of the plactic monoid for a semisimple Lie algebra \mathfrak{g} , we have $k = l$ and $c_i = c'_i$, for all $i = 1, \dots, k$. Thus $T = T'$. Since the 2-polygraph $\text{Path}(n)$ is terminating, and rewriting any non-standard tableau must terminate with a unique normal form, $\text{Path}(n)$ is confluent. \square

Every semisimple Lie algebra \mathfrak{g} admits a finite number of fundamental weights, then there is a finite number of L-S paths of shape ω_i , for $i = 1, \dots, n$. Thus the 2-polygraph $\text{Path}(n)$ is finite. Hence, by Lemmas 3.2.3 and 3.2.4, we obtain the following theorem.

3.2.5. Theorem. *For any semisimple Lie algebra \mathfrak{g} , the 2-polygraph $\text{Path}(n)$ is a finite convergent presentation of the plactic monoid for \mathfrak{g} .*

3.3. Finiteness properties of plactic monoids

A monoid is of *finite derivation type* (FDT_3) if it admits a finite presentation whose relations among the relations are finitely generated, see [SOK94]. The property FDT_3 is a natural extension of the properties of being finitely generated (FDT_1) and finitely presented (FDT_2). Using the notion of polygraphic resolution, one can define the higher-dimensional finite derivation type properties FDT_∞ , see [GM12]. They generalise in any dimension the finite derivation type FDT_3 . A monoid is said to be FDT_∞ if it admits a finite polygraphic resolution. By Corollary 4.5.4 in [GM12], a monoid with a finite convergent presentation is FDT_∞ . Then by Theorem 3.2.5, we have

3.3.1. Proposition. *For any semisimple Lie algebra \mathfrak{g} , plactic monoids for \mathfrak{g} satisfy the homotopical finiteness condition FDT_∞ .*

In the homological way, a monoid M is of *homological type* FP_∞ when there exists a resolution of M by projective, finitely generated $\mathbb{Z}M$ -modules. By Corollary 5.4.4 in [GM12] the property FDT_∞ implies the property FP_∞ . Hence we have

3.3.2. Proposition. *For any semisimple Lie algebra \mathfrak{g} , plactic monoids for \mathfrak{g} satisfy the homological finiteness property type FP_∞ .*

REFERENCES

- [Bak00] T. H. Baker. An insertion scheme for C_n crystals. In *Physical combinatorics (Kyoto, 1999)*, volume 191 of *Progr. Math.*, pages 1–48. Birkhäuser Boston, Boston, MA, 2000.
- [BCCL15] Leonid A. Bokut, Yuqun Chen, Weiping Chen, and Jing Li. New approaches to plactic monoid via Gröbner–Shirshov bases. *J. Algebra*, 423:301–317, 2015.
- [Bou68] Nicolas Bourbaki. *Éléments de mathématique. Fasc. XXXIV. Groupes et algèbres de Lie. Chapitre IV: Groupes de Coxeter et systèmes de Tits. Chapitre V: Groupes engendrés par des réflexions. Chapitre VI: systèmes de racines.* Actualités Scientifiques et Industrielles, No. 1337. Hermann, Paris, 1968.
- [CGM14] Alan J. Cain, Robert D. Gray, and António Malheiro. Crystal bases, finite complete rewriting systems, and biautomatic structures for Plactic monoids of types A_n , B_n , C_n , D_n , and G_2 . *ArXiv e-prints*, December 2014.
- [CGM15] Alan J. Cain, Robert D. Gray, and António Malheiro. Finite Gröbner–Shirshov bases for Plactic algebras and biautomatic structures for Plactic monoids. *J. Algebra*, 423:37–53, 2015.
- [DJM90] Etsurō Date, Michio Jimbo, and Tetsuji Miwa. Representations of $U_q(\mathfrak{gl}(n, \mathbb{C}))$ at $q = 0$ and the Robinson–Shensted [Schensted] correspondence. In *Physics and mathematics of strings*, pages 185–211. World Sci. Publ., Teaneck, NJ, 1990.
- [Ful97] William Fulton. *Young tableaux*, volume 35 of *London Mathematical Society Student Texts*. Cambridge University Press, Cambridge, 1997. With applications to representation theory and geometry.
- [GL12] Stéphane Gaussent and Peter Littelmann. One-skeleton galleries, the path model, and a generalization of Macdonald’s formula for Hall–Littlewood polynomials. *Int. Math. Res. Not. IMRN*, (12):2649–2707, 2012.
- [GM12] Yves Guiraud and Philippe Malbos. Higher-dimensional normalisation strategies for acyclicity. *Adv. Math.*, 231(3-4):2294–2351, 2012.
- [GM14] Yves Guiraud and Philippe Malbos. Polygraphs of finite derivation type. submitted, arXiv:1402.2587, 2014.
- [Hag14] Nohra Hage. Finite convergent presentation of plactic monoid of type C. arXiv:1412.0539, to appear in *International journal of algebra and computation*, 2014.
- [HM15] Nohra Hage and Philippe Malbos. Knuth’s coherent presentations of plactic monoid of type A. in preparation, 2015.
- [Hum78] James E. Humphreys. *Introduction to Lie algebras and representation theory*, volume 9 of *Graduate Texts in Mathematics*. Springer-Verlag, New York-Berlin, 1978. Second printing, revised.
- [Kas91] Masaki Kashiwara. Crystallizing the q -analogue of universal enveloping algebras. In *Proceedings of the International Congress of Mathematicians, Vol. I, II (Kyoto, 1990)*, pages 791–797. Math. Soc. Japan, Tokyo, 1991.

REFERENCES

- [Kas95] Masaki Kashiwara. On crystal bases. In *Representations of groups (Banff, AB, 1994)*, volume 16 of *CMS Conf. Proc.*, pages 155–197. Amer. Math. Soc., Providence, RI, 1995.
- [KN94] Masaki Kashiwara and Toshiki Nakashima. Crystal graphs for representations of the q -analogue of classical Lie algebras. *J. Algebra*, 165(2):295–345, 1994.
- [KO14] Łukasz Kubat and Jan Okniński. Gröbner-Shirshov bases for plactic algebras. *Algebra Colloq.*, 21(4):591–596, 2014.
- [Lec02] Cédric Lecouvey. Schensted-type correspondence, plactic monoid, and jeu de taquin for type C_n . *J. Algebra*, 247(2):295–331, 2002.
- [Lec06] Cédric Lecouvey. Combinatorics of crystal graphs and Kostka-Foulkes polynomials for the root systems B_n , C_n and D_n . *European J. Combin.*, 27(4):526–557, 2006.
- [Lec07] Cédric Lecouvey. Combinatorics of crystal graphs for the root systems of types A_n , B_n , C_n , D_n and G_2 . In *Combinatorial aspect of integrable systems*, volume 17 of *MSJ Mem.*, pages 11–41. Math. Soc. Japan, Tokyo, 2007.
- [Lit94] Peter Littelmann. A Littlewood-Richardson rule for symmetrizable Kac-Moody algebras. *Invent. Math.*, 116(1-3):329–346, 1994.
- [Lit95] Peter Littelmann. Paths and root operators in representation theory. *Ann. of Math. (2)*, 142(3):499–525, 1995.
- [Lit96] Peter Littelmann. A plactic algebra for semisimple Lie algebras. *Adv. Math.*, 124(2):312–331, 1996.
- [LLT95] Alain Lascoux, Bernard Leclerc, and Jean-Yves Thibon. Crystal graphs and q -analogues of weight multiplicities for the root system A_n . *Lett. Math. Phys.*, 35(4):359–374, 1995.
- [Lop14] Viktor Lopatkin. Cohomology Rings of Plactic Monoid Algebra via Gröbner-Shirshov Basis. *ArXiv e-prints*, November 2014.
- [Lot02] M. Lothaire. *Algebraic combinatorics on words*, volume 90 of *Encyclopedia of Mathematics and its Applications*. Cambridge University Press, Cambridge, 2002.
- [LS81] Alain Lascoux and Marcel-P. Schützenberger. Le monoïde plaxique. In *Noncommutative structures in algebra and geometric combinatorics (Naples, 1978)*, volume 109 of *Quad. "Ricerca Sci."*, pages 129–156. CNR, Rome, 1981.
- [SOK94] Craig C. Squier, Friedrich Otto, and Yuji Kobayashi. A finiteness condition for rewriting systems. *Theoret. Comput. Sci.*, 131(2):271–294, 1994.

NOHRA HAGE
Université de Lyon,
Institut Camille Jordan, CNRS UMR 5208
Université Jean Monnet
23, boulevard du docteur Paul Michelon,
42000 Saint-Etienne cedex, France.
nohra.hage@univ-st-etienne.fr